



(12) PATENT ABRIDGMENT (11) Document No. AU-B-21926/95
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 698145

- (54) Title
SYSTEM FOR PH-NEUTRAL LONGLIFE PRECAST ELECTROPHORESIS GEL
- International Patent Classification(s)
(51)⁶ **G01N 027/447**
- (21) Application No. : **21926/95** (22) Application Date : **29.03.95**
- (87) PCT Publication Number : **WO95/27197**
- (30) Priority Data
- (31) Number (32) Date (33) Country
08/221939 31.03.94 US UNITED STATES OF AMERICA
- (43) Publication Date : **23.10.95**
- (44) Publication Date of Accepted Application : **22.10.98**
- (71) Applicant(s)
NOVEL EXPERIMENTAL TECHNOLOGY
- (72) Inventor(s)
SHELDON ENGELHORN; TIMOTHY V UPDYKE
- (74) Attorney or Agent
CULLEN & CO , GPO Box 1074, BRISBANE QLD 4001
- (56) Prior Art Documents
EP 566784
US 4481094
US 4209373
- (57) Claim

1. A system for gel electrophoresis comprising:
an electrophoresis gel having two ends; said gel being saturated with a gel buffer solution;
one end of the gel being in contact with an anode buffer solution;
the other end of the gel being in contact with a cathode buffer solution;
said gel buffer solution comprising a monovalent organic amine or substituted amine with a pK_a near neutrality, titrated with hydrochloric acid (HCl) to a substantially neutral pH;
said cathode buffer comprising a solution of a zwitterionic buffer selected from the group of 3-(N-morpholino)-propanesulfonic acid (MOPS), 2-(N-morpholino)-ethanesulfonic acid (MES) N-(2-acetamido)-2-aminoethanesulfonic acid (ACES), 2-(N-morpholino)-2-hydroxypropanesulfonic acid (MOPSO), N-tris-(hydroxymethyl)-2-ethanesulfonic acid (TES), N-2-hydroxyethyl-piperazine-N-2-ethanesulfonic

acid (HEPES), or 3-(N-tris-(hydroxymethyl) methylamino)-2-hydroxypropanesulfonic acid (TAPSO), titrated to a substantially neutral pH with sodium hydroxide or an organic base.

13. A system for gel electrophoresis comprising:
 - a precast polyacrylamide electrophoresis gel having two ends;
 - wherein said gel is saturated with a gel buffer solution;
 - wherein one end of the gel is in contact with an anode buffer solution;
 - wherein the other end of the gel is in contact with a cathode buffer solution;
 - wherein said gel buffer solution comprises a monovalent organic amine or substituted amine with a pK_a near neutrality, titrated with hydrochloric acid (HCl) to a substantially neutral pH; and
 - wherein said cathode buffer comprises sulfite.

OPI DATE 23/10/95 APPLN. ID 21926/95
AOJP DATE 07/12/95 PCT NUMBER PCT/US95/03647



AU9521926

(PCT)

(51) International Patent Classification ⁶ :

G01N 27/447

A1

(11) International Publication Number:

WO 95/27197

(43) International Publication Date:

12 October 1995 (12.10.95)

(21) International Application Number: PCT/US95/03647

(22) International Filing Date: 29 March 1995 (29.03.95)

(30) Priority Data:

08/221,939

31 March 1994 (31.03.94)

US

(71) Applicant: NOVEL EXPERIMENTAL TECHNOLOGY

[US/US]; ~~4202 Sorrento Valley Boulevard, San Diego, CA~~

~~92121 (US)~~

San Diego CA 92121 United States of America

(72) Inventors: ENGELHORN, Sheldon; 734 Oak Burl Lane, Encinitas, CA 92024 (US). UPDYKE, Timothy, V.; 37349 Avenida Chapala, Temecula, CA 92592 (US).

(74) Agents: BEAMER, Norman, H. et al.; Fish & Neave, 1251 Avenue of the Americas, New York, NY 10020 (US).

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).

Published

With international search report.

America



(54) Title: SYSTEM FOR PH-NEUTRAL LONGLIFE PRECAST ELECTROPHORESIS GEL

(57) Abstract

A gel and buffer system for gel electrophoresis wherein separation occurs at neutral pH and proteins remain completely reduced.

SYSTEM FOR PH-NEUTRAL LONGLIFE
PRECAST ELECTROPHORESIS GEL

This invention relates to techniques for gel
5 electrophoresis. More particularly this invention
relates to a novel system for gel electrophoresis at
approximately neutral pH.

Background Of the Invention

Gel electrophoresis is a common procedure for
10 the separation of biological molecules, such as DNA,
RNA, polypeptides and proteins. In gel
electrophoresis, the molecules are separated into bands
according to the rate at which an imposed electric
field causes them to migrate through a filtering gel.

15 The basic apparatus used in this technique
consists of a gel enclosed in a glass tube or
sandwiched as a slab between glass or plastic plates.
The gel has an open molecular network structure,
defining pores which are saturated with an electrically
20 conductive buffered solution of a salt. These pores
through the gel are large enough to admit passage of
the migrating macromolecules.

The gel is placed in a chamber in contact
with buffer solutions which make electrical contact
25 between the gel and the cathode or anode of an
electrical power supply. A sample containing the

- 2 -

macromolecules and a tracking dye is placed on top of the gel. An electric potential is applied to the gel causing the sample macromolecules and tracking dye to migrate toward the bottom of the gel. The

5 electrophoresis is halted just before the tracking dye reaches the end of the gel. The locations of the bands of separated macromolecules are then determined. By comparing the distance moved by particular bands in comparison to the tracking dye and macromolecules of

10 known mobility, the mobility of other macromolecules can be determined. The size of the macromolecule can then be calculated.

The rate of migration of macromolecules through the gel depends upon three principle factors:

15 the porosity of the gel; the size and shape of the macromolecule; and the charge density of the macromolecule. It is critical to an effective electrophoresis system that these three factors be precisely controlled and reproducible from gel to gel

20 and from sample to sample. However, maintaining uniformity between gels is difficult because each of these factors is sensitive to many variables in the chemistry of the gel system.

Polyacrylamide gels are commonly used for

25 electrophoresis. Polyacrylamide gel electrophoresis or PAGE is popular because the gels are optically transparent, electrically neutral and can be made with a range of pore sizes. The porosity of a polyacrylamide gel is in part defined by the total

30 percentage of acrylamide monomer plus crosslinker monomer ("%T") it contains. The greater the concentration, the less space there is between strands of the polyacrylamide matrix and hence the smaller the pores through the gel. An 8% polyacrylamide gel has

35 larger pores than a 12% polyacrylamide gel. An 8%

- 3 -

polyacrylamide gel consequently permits faster migration of macromolecules with a given shape, size and charge density. When smaller macromolecules are to be separated, it is generally preferable to use a gel
5 with a smaller pore size such as a 20% gel. Conversely for separation of larger macromolecules, a gel with a larger pore size is often used, such as an 8% gel.

Pore size is also dependent upon the amount of crosslinker used to polymerize the gel. At any
10 given total monomer concentration, the minimum pore size for a polyacrylamide gel is obtained when the ratio of total monomer to crosslinker is about 20:1, (the usual expression for this ratio would be "5%C").

Several factors may cause undesirable
15 variation in the pore size of gels. Pore size can be increased by incomplete gel polymerization during manufacture. Hydrolysis of the polyacrylamide after polymerization can create fixed negative charges and break down the crosslinks in the gel, which will
20 degrade the separation and increase the pore size. An ideal gel system should have a reproducible pore size and no fixed charge (or at least a constant amount) and should be resistant to change in chemical characteristics or the pore size due to hydrolysis.

25 The size of the macromolecule varies between different macromolecules; the smaller and more compact the macromolecule the easier it will be for the macromolecule to move through the pores of a given gel. Given a constant charge density, the rate of migration
30 of a macromolecule is inversely proportional to the logarithm of its size.

For accurate and reproducible electrophoresis, a given type of macromol cule should preferably take on a single form in the gel. One
35 difficulty with maintaining uniformity of the shape of

- 4 -

proteins during gel electrophoreses is that disulfide bonds can be formed by oxidation of pairs of cysteine amino acids. Different oxidized forms of the protein then have different shapes and, therefore, migrate
5 through the gel run with slightly different mobilities (usually faster than a completely reduced protein, since the maximum stokes radius and minimum mobility should occur with a completely unfolded form). A heterogeneous mixture of forms leads to apparent band
10 broadening. In order to prevent the formation of disulfide bonds, a reducing agent such as dithiothreitol (DTT) is usually added to the samples to be run.

The charge density of the migrating molecule
15 is the third factor affecting its rate of migration through the gel -- the higher the charge density, the more force will be imposed by the electric field upon the macromolecule and the faster the migration rate subject to the limits of size and shape. In SDS-PAGE
20 electrophoresis the charge density of the macromolecules is controlled by adding sodium dodecyl sulfate ("SDS") to the system. SDS molecules associate with the macromolecules and impart a uniform charge density to them substantially negating the effects of
25 any innate molecular charge.

SDS PAGE gels are usually poured and run at basic pH. The most common PAGE buffer system employed for the separation of proteins is that developed by Ornstein (1) and modified for use with SDS by Laemmli
30 (2). Laemmli, U.K. (1970) Nature 227, 680-686. The Laemmli buffer system consists of 0.375 M Tris (tris (hydroxy methyl) amino-methane), titrated to pH 8.8. with HCl, in the separating gel. The stacking gel consists of 0.125 M Tris, titrated to pH 6.8. The
35 anode and cathode running buffers contain 0.024 M Tris,

- 5 -

0.192 M glycine, 0.1% SDS. An alternative buffer system is disclosed by Schaeffer and von Jagow. Schagger, H. and von Jagow, G., Anal. Biochem. 1987, 166, 368-379. The stacking gel contains 0.75 M Tris, 5 titrated to pH 8.45 with HCl. The separating gel contains 0.9 M Tris, titrated to pH 8.45 with HCl. The cathode buffer contains 0.1 M Tris, 0.1M Tricine, 0.1% SDS. The anode buffer contains 0.2 M Tris, titrated to pH 8.9 with HCl. For both of these systems Tris is the 10 "common ion" which is present in the gel and in the anode and cathode buffers.

In the Laemmli system, the pH of the trailing phase in the stacking gel is about 8.9. In the separating gel, the trailing phase pH is about 9.7. At 15 this pH, primary amino groups of proteins react readily with unpolymerized acrylamide, thiol groups are more subject to oxidation to disulfides, or reaction with unpolymerized polyacrylamide, than at neutral pH and acrylamide itself is subject to hydrolysis.

20 The need for uniformity and predictability is magnified in precast electrophoresis gels which are manufactured by an outside vendor and then shipped to the laboratory where the electrophoresis will be performed. Precast gels must control the properties 25 discussed above and they must be able to maintain this control throughout shipping and storage. The shelf life of many precast gels is limited by the potential for hydrolysis of acrylamide during storage at the high pH of the gel buffer.

30 It is a disadvantage of a high pH gel that the polyacrylamide gel is subject to degradation by hydrolysis and has a limited shelf-life.

It is a further disadvantage of a high pH gel that proteins react readily with unpolymerized

- 6 -

acrylamide which may interfere with subsequent analysis of the protein such as peptide sequencing.

It is a still further disadvantage of a high pH gel that thiol groups are subject to oxidation to
5 disulfides causing a decreased resolution of separated macromolecules.

Summary of the Invention

It is an object of this invention to produce a neutral gel system that reduces protein reaction with
10 unpolymerized acrylamide thereby enhancing yield and resolution.

It is a further object of this invention to produce a neutral gel system that prevents formation of disulfides from free thiol groups thereby enhancing
15 yield and resolution.

It is a still further object of this invention to produce a neutral gel system that reduces degradation of the polyacrylamide gel by hydrolysis thereby increasing the useful shelf-life of a precast
20 gel.

In accordance with this invention, applicants describe a gel and buffer system wherein separation occurs at neutral pH and proteins remain completely reduced. The above and other objects and advantages of
25 the present invention will be apparent upon consideration of the following detailed description.

Detailed Description Of The Invention

Applicants describe a gel and buffer system wherein separation occurs at neutral pH and proteins
30 remain completely reduced. Advantageously, at this neutral pH, primary amino groups of proteins react less readily with unpolymerized acrylamide because protonation of protein amino groups greatly reduces

their reactivity to acrylamide or other related blocking agents. Furthermore, at this neutral pH, thiol groups are less subject to oxidation than at higher pH and polyacrylamide itself is less subject to hydrolysis.

The result is a gel system with improved stability of the gel matrix and stock solutions. Gels prepared according to this system can be stored under refrigeration for over a year without loss of performance due to acrylamide hydrolysis. Also, stock buffers without reducing agents and stock gel solutions without polymerization initiator can be stored for at least several weeks at room temperature with no loss of performance. An additional benefit is that a single gel recipe, using the same buffer for the stacking and separating gels, can be used with two different running buffers to give two separation systems. Using this feature, an 8% gel, for example, can cover a protein separation range of 2 to 200 kDa.

In one embodiment of this invention a polyacrylamide gel of between about 3% and about 25% (%T) acrylamide is polymerized using from about 1% to about 6% crosslinker (%C) using a gel buffer comprising a primary organic amine or substituted amine with a pK_a near neutrality, titrated with approximately half as much HCl (on a molar basis), so that the pH of the buffer is approximately neutral. In a preferred embodiment the gel is polymerized using from about 2% to about 5% crosslinker (%C) using a gel buffer comprising Bis-Tris [Bis-(2-hydroxyethyl) iminotris (hydroxymethyl) methane] titrated with HCl. Different separation characteristics can be obtained by running the gel with either a MOPS (3-[N-Morpholino] propanesulfonic acid) or MES (2-(N-Morpholino) ethanesulfonic acid), buffer. 2mM to 10 mM TGA or 2mM



to 10mM sodium bisulfite is added to the running buffer to maintain a reducing environment in the gel during electrophoresis. This and other embodiments can be understood by reference to the following illustrative and comparative examples.

5

Examples

Bis-Tris, MES, and MOPS were purchased from Sigma (St. Louis, MO) or Research Organics (Cleveland, Ohio). Thioglycolic acid (TGA), DTT and mercaptoethanol (BME) were from Sigma. All other chemicals were reagent, "ultra pure" or "electrophoresis grade" from standard sources. Gels were cast in 1mm thickness mini-gel cassettes from Novex (San Diego CA) and run in Novex minicell.

The Bis-Tris separating gel and stacking gels were prepared from a 30%T/2.5%C acrylamide/BIS stock solution and a 7X Bis-Tris stock solution (2.5M Bis-Tris, 1.5M HCl, pH 6.5). To prepare the separating gel, the stock solutions were blended with ultra pure water to a final concentration 8%T, 0.357M Bis-Tris, to which was added 0.2 ul/ml N,N,N',N'-tetra-methylene-diamine (TEMED). After degassing, 2.0ul/ml of a 10% solution of APS was added, the gel was immediately poured into the cassette then overlaid with water. Polymerization was allowed to proceed for at least 30 minutes at room temperature, the water was removed and a 4% stacking gel applied. The stacking gel was prepared in the same fashion as the separating gel, except that the final concentration obtained was 4%T, temed concentration was increased to 0.4ul/ml and the APS increased to 5.0ul/ml.

MOPS running buffer consisted of 50mM MOPS, 50mM Bis-Tris (or TRIS), 0.1% SDS, 1mM EDTA. MES running buffer consisted of 50mM MES, 50mM Bis-Tris (or TRIS), 0.1% SDS, 1mM EDTA. Sample buffer (2X)



- 9 -

consisted of 0.25M Bis-Tris, 0.15M HCl, 10% (w/v) Glycerol, 2% SDS, 1mM EDTA, 0.03% Serva Blue G, and 200mM DTT. Samples containing a set of protein standards were heated for 15 min at 70 degrees before application. Bovine serum albumin (BSA), chicken egg ovalbumin, alkylated insulin A and B chain, soybean trypsin inhibitor, and bovine erythrocyte carbonic anhydrase were included in the standard. Sample volume was 5ul in all cases.

10

Example 1

The protein standards were separated on an 8% Bis-Tris gel with MOPS running buffer in the absence of a reducing agent. The resulting separation pattern was very similar to that obtained on an 8% TRIS/glycine gel (Laemmli), with proteins 20,000 and smaller remaining in the stack along with the tracking dye. The BSA band was somewhat diffuse and shifted toward the anode. The Ovalbumin band was also somewhat diffuse.

20

Example 2

The protein standards were separated on an 8% Bis-Tris gel with MOPS running buffer in the presence of TGA in the cathode buffer. Again, the separation pattern was very similar to that obtained on an 8% TRIS/glycine (Laemmli) gel, with proteins 20,000 and smaller remaining in the stack along with the tracking dye. The presence of the reducing agent, 5mM TGA, in the cathode buffer provided for better resolution of the proteins BSA and Ovalbumin compared to the gel run without TGA.

30

Example 3

The protein standards were separated on an 8% Bis-Tris gel with MOPS running buffer in the presence

- 10 -

of sodium bisulfite in the cathode buffer. Again, the separation pattern was very similar to that obtained on an 8% TRIS/glycine (Laemmli) gel, with proteins 20,000 and smaller remaining in the stack along with the tracking dye. The presence of the reducing agent, 5mM sodium bisulfite, in the cathode buffer provided for better resolution of the proteins BSA and Ovalbumin compared to the gel run without sodium bisulfite.

Example 4

The protein standards were separated on an 8% Bis-Tris gel with MES running buffer in the absence of a reducing agent. The protein separation was very similar to that obtained from a 12% TRIS/tricine (Schaeffer) gel. All proteins were resolved from the stack including insulin A and B chain (3500 and 2500 daltons, respectively). When the gel is run without TGA, soybean trypsin inhibitor had a more prominent doublet.

Example 5

The protein standards were separated on an 8% Bis-Tris/Cl gel with Bis-Tris/MES running buffer in the presence of TGA in the cathode buffer. Again, all proteins were resolved from the stack including insulin A and B chain (3500 and 2500 daltons, respectively). The presence of the reducing agent, 5mM TGA, in the cathode buffer provided for better resolution of the protein soybean trypsin inhibitor. Carbonic anhydrase ran as a tight, sharp band under all conditions tested.

Example 6

The protein standards were separated on an 8% Bis-Tris/Cl gel with Bis-Tris/MES running buffer in the presence of sodium bisulfite in the cathode buffer. Again, all proteins were resolved from the stack

- 11 -

including insulin A and B chain (3500 and 2500 daltons, respectively). The presence of the reducing agent, 5mM sodium bisulfite, in the cathode buffer provided for better resolution of the protein soybean trypsin inhibitor. Carbonic anhydrase ran as a tight, sharp band under all conditions tested.

Although MES and MOPS were selected as desirable running buffers because the resulting system has separation characteristics similar to the commonly used Laemmli and Schaeffer gel systems, it was found that a range of buffers are suitable for use in this system. Among the additional buffers giving good results were ACES ([N-(2-Acetomido)]-2-aminoethanesulfonic acid, MOPSO (2-[N-Morpholino]-2-hydroxypropanesulfonic acid, TES (N-Tris-(hydroxymethyl)-2-ethanesulfonic acid, BES (N,N-BIS-(Hydroxyethyl)-2-aminoethanesulfonic acid, HEPES (N-2-Hydroxyethyl-piperazine-n-2-ethanesulfonic acid), TAPSO (3-(N-tris-(Hydroxymethyl) methylamino)-2-hydroxypropanesulfonic acid.

All the proteins that exhibit some band broadening and/or mobility shifts when run in the absence of TGA or sodium bisulfite, have in common a composition that includes multiple cysteines (BSA, for instance, has 35 cysteines). On the other hand carbonic anhydrase, which always runs cleanly, has no cysteines. Moreover, if the reduced proteins are alkylated before running, they run as sharp homogeneous bands even in the absence of a reducing agent.

Cysteine-containing proteins appear to give generally sharper bands in the Laemmli system than the neutral system, when both are run with 100mM mercaptoethanol or DTT in the sample buffer but without TGA in the running buffer. Since thiol oxidation is more favored as the pH increases, it would be expected

- 12 -

that the higher pH of the Laemmli system would cause oxidation of disulfide to be at least as pronounced as it is in the neutral pH system. However, DTT and similar "neutral" thiol reducing agents are weak acids
5 (with pKa's around pH 8-9). Thus, at basic pH, these reducing agents migrate into the gel and, if present at sufficient concentration, provide some protection against oxidation of sulfhydryls. At a neutral separating pH, DTT from the sample buffer is in an
10 uncharged form and will remain behind in the sample well. Thus, no reducing agent migrates into the gel.

To maintain proteins in a reduced form during electrophoresis at neutral pH, it was found advantageous to use a reducing agent that would migrate
15 into the gel at neutral pH. Sodium Bisulfite (2-10mM) was found to maintain a reducing environment in the gel during electrophoresis. Fully reduced TGA (or similar negatively charged thiols) give similar results at comparable concentrations. However, partially oxidized
20 TGA will promote partial oxidation of protein thiols. Because reduction (oxidation) of protein thiols will take place via disulfide interchange, the ratio of reduced to oxidized thiols in the protein will substantially reflect the ratio of reduced to oxidized
25 thiols in the TGA. Conversely, sulfite oxidizes to sulfate, which does not participate in redox reactions under conditions found in the gel. Therefore, regardless of the sulfite/sulfate ratio in a partially oxidized preparation of sulfite, as long as sufficient
30 sulfite remains, proteins will be protected against thiol oxidation.

It was also found that TRIS could be substituted for Bis-Tris in the running buffer with no visible effect on the separation quality. Bis-Tris may
35 be preferred where the protein will be intentionally

- 13 -

modified post-separation. Bis-Tris is a tertiary amine and will not interfere with the protein modifying agents which react through primary amines. TRIS, however is the preferred choice for routine use,
5 because it is available at significantly lower cost than Bis-Tris.

While the invention has been explained in relation to its preferred embodiments, it is to be understood that various modifications thereof will
10 become apparent to those skilled in the art. The foregoing disclosure is not intended or to be construed to limit the present invention, or to otherwise exclude any such other embodiments, adaptations, variations and equivalent arrangements, the present invention being
15 limited only by the claims appended hereto and the equivalents thereof.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A system for gel electrophoresis comprising:
an electrophoresis gel having two ends; said gel being saturated
5 with a gel buffer solution;
one end of the gel being in contact with an anode buffer
solution;

the other end of the gel being in contact with a cathode buffer
solution;

10 said gel buffer solution comprising a monovalent organic amine
or substituted amine with a pK_a near neutrality, titrated with hydrochloric acid
(HCl) to a substantially neutral pH;

said cathode buffer comprising a solution of a zwitterionic buffer
selected from the group of 3-(N-morpholino)-propanesulfonic acid (MOPS), 2-
15 (N-morpholino)-ethanesulfonic acid (MES) N-(2-acetamido)-2-
aminoethanesulfonic acid (ACES), 2-(N-morpholino)-2-
hydroxypropanesulfonic acid (MOPSO), N-tris-(hydroxymethyl)-2-
ethanesulfonic acid (TES), N-2-hydroxyethyl-piperazine-N-2-ethanesulfonic
acid (HEPES), or 3-(N-tris-(hydroxymethyl) methylamino)-2-
20 hydroxypropanesulfonic acid (TAPSO), titrated to a substantially neutral pH
with sodium hydroxide or an organic base.

2. The system of claim 1 wherein the monovalent organic amine is
bis-(2-hydroxyethyl) iminotris (hydroxymethyl) methane (Bis-Tris).

25 3. The system of claim 2 wherein the gel is a polyacrylamide gel.

4. The system of claim 3 wherein the cathode buffer solution
comprises sulfite.

30 5. The system of claim 3 wherein the cathode buffer solution
comprises a negatively charged thiol.



6. The system of claim 3 wherein the cathode buffer solution comprises thioglycolic acid (TGA).

7. The system of claim 3 wherein the anode buffer comprises tris
5 (hydroxy methyl) amino-methane (TRIS).

8. A system for gel electrophoresis comprising:
a precast polyacrylamide electrophoresis gel having two ends;
said gel being saturated with a gel buffer solution;
10 one end of the gel being in contact with an anode buffer
solution;
the other end of the gel being in contact with a cathode buffer
solution;
said gel buffer solution comprising Bis-Tris titrated with
15 hydrochloric acid (HCl) to a substantially neutral pH;
said cathode buffer comprising a solution of a zwitterionic buffer
selected from the group of 3-(N-morpholino)-propanesulfonic acid (MOPS), 2-
(N-morpholino)-ethanesulfonic acid (MES), N-(2-acetamido)-2-
aminoethanesulfonic acid (ACES), 2-(N-morpholino)-2-
20 hydroxypropanesulfonic acid (MOPSO), N-tris-(hydroxymethyl)-2-
ethanesulfonic acid (TES), N-2-hydroxyethyl-piperazine-N-2-ethanesulfonic
acid (HEPES), or 3-(N-tris-(hydroxymethyl) methylamino)-2-
hydroxypropanesulfonic acid (TAPSO), titrated to a substantially neutral pH
with sodium hydroxide or an organic base.

25

9. The system of claim 8 wherein the anode buffer comprises tris
(hydroxy methyl) amino-methane (TRIS).

10. The system of claim 8 wherein the cathode buffer comprises
30 sulfite.

11. The system of claim 8 wherein the cathode buffer comprises a
negatively charged thiol.



12. The system of claim 8 wherein the cathode buffer comprises thioglycolic acid (TGA).

- 5 13. A system for gel electrophoresis comprising:
a precast polyacrylamide electrophoresis gel having two ends;
wherein said gel is saturated with a gel buffer solution;
wherein one end of the gel is in contact with an anode buffer
solution;
10 wherein the other end of the gel is in contact with a cathode
buffer solution;
wherein said gel buffer solution comprises a monovalent organic
amine or substituted amine with a pK_a near neutrality, titrated with
hydrochloric acid (HCl) to a substantially neutral pH; and
15 wherein said cathode buffer comprises sulfite.

DATED this 11th day of September 1998
NOVEL EXPERIMENTAL TECHNOLOGY

20

By their Patent Attorneys
CULLEN & CO.



INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 95/03647

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 G01N27/447

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 566 784 (HYMO CORP.) 27 October 1993 see page 3, line 23 - page 4, line 13 ---	1
Y	US,A,4 481 094 (A. FERNANDEZ DE CASTRO) 6 November 1984 see column 3, line 33 - column 4, line 49 ---	1
A	US,A,4 209 373 (B. I. BLUESTEIN) 24 June 1980 see abstract ---	1
A	US,A,3 948 743 (J. F. MONTHONY) 6 April 1976 see column 3, line 51 - line 55 ---	1
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

10 July 1995

Date of mailing of the international search report

24.07.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Duchatellier, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/03647

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,5 275 708 (R. E. AKINS) 4 January 1994 see claim 3 ---	1
A	EP,A,0 518 475 (BECKMAN INSTRUMENTS, INC.) 16 December 1992 see abstract -----	1

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/03647

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0566784	27-10-93	JP-A- 4184163	01-07-92
US-A-4481094	06-11-84	NONE	
US-A-4209373	24-06-80	NONE	
US-A-3948743	06-04-76	NONE	
US-A-5275708	04-01-94	NONE	
EP-A-0518475	16-12-92	US-A- 5120413	09-06-92
		AU-A- 1628792	03-12-92